

MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS-196 $^{\circ}$ A

STAL MANNEY OF SECTIONS AND SECTION OF SECTI

CURITY CLASSIFICATION OF THIS PAGE				
OTHE FILE CUPPERORT DOCU	MENTATION	PAGE		
REPORT SECURITY CLASSIFICATION	16. RESTRICTIVE			
DTIC				
AFI FCTE		/AVAILABILITY O		
AD-A190 042 FEB 2 9 1988	2	for Public		
FEB 2 9 1988 RE		cion Unlimit		E B/CS
QQ LI	AFC	**************************************	88-0	
NAME OF PERFORMING ORGANIZATION 6b. OFFICE SYMBOL (If applicable)	1	ONITORING ORGA	NIZATION	
North Dakota State Univ.	AFOSR/N			
ADDRESS (City, State, and ZIP Code)	7b. ADDRESS (C)	ty, State, and ZIP	Code)	
Fargo, North Dakota 58105	Buildin	g 410		
	Bolling	AFB 20332		
NAME OF FUNDING/SPONSORING ORGANIZATION 8b. OFFICE SYMBOL (If applicable)	9. PROCUREMEN	T INSTRUMENT ID	ENTIFICATION	NUMBER
AFOSR NC	arosi2 84			
ADDRESS (City, State, and ZIP Code)	10. SOURCE OF	FUNDING NUMBER	TASK	WORK UNIT
Building 410	ELEMENT NO.	NO.	NO.	ACCESSION N
Bolling AFB 20322 TITLE (Include Security Classification)	61102F	2303	B2	
Philip Boudjouk TYPE OF REPORT Final Philip Boudjouk 13b. TIME COVERED FROM 11/1/83 TO 10/31/8	14. DATE OF REPO			IGE COUNT
Philip Boudjouk Type of Report 13b. TIME COVERED	14. DATE OF REPO	ORT (Year, Month,		
Philip Boudjouk TYPE OF REPORT Final SUPPLEMENTARY NOTATION Philip Boudjouk 13b. TIME COVERED FROM 11/1/83 TO 10/31/8	14. DATE OF REPO	ORT (Year, Month, 88, 1, 5	5 2	1
Philip Boudjouk TYPE OF REPORT Final SUPPLEMENTARY NOTATION COSATI CODES 18. SUBJECT TERMS	14. DATE OF REPO	ORT (Year, Month, 88, 1, 5	od identify by	block number)
Philip Boudjouk TYPE OF REPORT Final SUPPLEMENTARY NOTATION COSATI CODES FIELD GROUP Philip Boudjouk 13b. TIME COVERED FROM 11/1/83 TO 10/31/8 18 SUBJECT TERMS Ultraso	14. DATE OF REPO	ORT (Year, Month, 88, 1, 5) se if necessary and nistry, Sili	od identify by	block number) nosilanes,
Philip Boudjouk TYPE OF REPORT Final SUPPLEMENTARY NOTATION COSATI CODES FIELD GROUP SUB-GROUP Ultraso Organos	(Continue on reversional, Sonochemelenium, Hydro	ORT (Year, Month, 88, 1, 5) se if necessary and nistry, Sili	od identify by	block number) nosilanes,
Philip Boudjouk a. TYPE OF REPORT Final SUPPLEMENTARY NOTATION COSATI CODES FIELD GROUP SUB-GROUP Ultraso Organos ABSTRACT (Continue on reverse if necessary and identify by block	(Continue on revenund, Sonochen elenium, Hydroumber)	DRT (Year, Month, 88, 1, 5) se if necessary and mistry, Silings of the control of	od identify by con, Organilylenes,	block number) nosilanes,
Philip Boudjouk Type of REPORT Final SUPPLEMENTARY NOTATION COSATI CODES FIELD GROUP SUB-GROUP Ultraso Organos ABSTRACT (Continue on reverse if necessary and identify by block The objectives of this program for the	(Continue on revenund, Sonochenelenium, Hydronumber)	DRT (Year, Month, 88, 1, 5) se if necessary and stry, Siling cosilation, S /84 - 10/31/	od identify by con, Organilylenes,	block number) nosilanes, Silacenium
Philip Boudjouk TYPE OF REPORT Final COSATI CODES FIELD GROUP SUB-GROUP The objectives of this program for the 1) to continue to extend our research on ul 2) to continue to develop the new chemistry	Continue on revenund, Sonocherelenium, Hydronumber) period 11/1 trasonically of low temp	per if necessary and istry, Silicosilation, S /84 - 10/31/ accelerated erature generature generature.	d identify by con, Organilylenes, /87 were d reaction	block number) nosilanes, Silacenium
Philip Boudjouk TYPE OF REPORT Final COSATI CODES FIELD GROUP SUB-GROUP The objectives of this program for the 1) to continue to extend our research on ul 2) to continue to develop the new chemistry silylenes containing large groups on the si	Continue on reversion of low templicon atom;	per if necessary and istry, Sili cosilation, S /84 - 10/31/ accelerated erature generally to develop	d identify by con, Organilylenes, 87 were d reaction erated op routes	block number) nosilanes, Silacenium
Philip Boudjouk TYPE OF REPORT Final COSATI CODES FIELD GROUP SUB-GROUP The objectives of this program for the objectives of this program for the l) to continue to extend our research on ul 2) to continue to develop the new chemistry silylenes containing large groups on the si silylenium ions in solution and investigate	(Continue on rever- und, Sonocher elenium, Hydr number) e period 11/1 trasonically of low temp licon atom;	per if necessary and mistry, Sili cosilation, S /84 - 10/31/ accelerated erature generature generature; 4) to development; 4) to	d identify by con, Organilylenes, 87 were d reaction erated op routes develop t	block number) nosilanes, Silacenium
Philip Boudjouk 13b. TIME COVERED FROM 11/1/83 TO 10/31/8 5. SUPPLEMENTARY NOTATION COSATI CODES FIELD GROUP SUB-GROUP Ultraso Organos ABSTRACT (Continue on reverse if necessary and identify by block The objectives of this program for the 1) to continue to extend our research on ul 2) to continue to develop the new chemistry silylenes containing large groups on the si silylenium ions in solution and investigate chemistry of spiropentasilane derivatives; syntheses of novel silicon-chalcogenide and	(Continue on rever- und, Sonocher elenium, Hydr number) period 11/1 trasonically of low temp licon atom; their chemi 5) to initia	DRT (Year, Month, 88, 1, 5) se if necessary and nistry, Silitosilation, S /84 - 10/31/ accelerated erature generature generature generature; 4) to te programs nsition meta	con, Organilylenes, 87 were reaction erated op routes develop to in the al	block number) nosilanes, Silacenium
Philip Boudjouk TYPE OF REPORT Final COSATI CODES FIELD GROUP SUB-GROUP The objectives of this program for the 1) to continue to extend our research on ul 2) to continue to extend our research on ul 2) to continue to develop the new chemistry silylenes containing large groups on the si silylenium ions in solution and investigate chemistry of spiropentasilane derivatives; syntheses of novel silicon-chalcogenide and compounds; and, 6) to investigate the elect	Continue on revenund, Sonochere elenium, Hydronically of low templicon atom; their chemis) to initial silicon-transchemistry	per if necessary and instry, Silinguistry, S	con, Organilylenes, /87 were interested op routes develop to in the all alized	block number) nosilanes, Silacenium
Philip Boudjouk 13b. TIME COVERED FROM 11/1/83 TO 10/31/8 SUPPLEMENTARY NOTATION COSATI CODES FIELD GROUP SUB-GROUP Ultraso Organos ABSTRACT (Continue on reverse if necessary and identify by block The objectives of this program for the 1) to continue to extend our research on ul 2) to continue to develop the new chemistry silvlenes containing large groups on the si silvlenium ions in solution and investigate chemistry of spiropentasilane derivatives; syntheses of novel silicon-chalcogenide and compounds; and, 6) to investigate the elect organosilanes. In this report we summarize	Continue on revenund, Sonochere elenium, Hydronically of low templicon atom; their chemis) to initial silicon-transchemistry	per if necessary and instry, Silinguistry, S	con, Organilylenes, /87 were interested op routes develop to in the all alized	block number) nosilanes, Silacenium as; to
Philip Boudjouk TYPE OF REPORT Final COSATI CODES FIELD GROUP SUB-GROUP The objectives of this program for the l) to continue to extend our research on ul 2) to continue to extend our research on ul 2) to continue to develop the new chemistry silylenes containing large groups on the si silylenium ions in solution and investigate chemistry of spiropentasilane derivatives; syntheses of novel silicon-chalcogenide and compounds; and, 6) to investigate the elect	Continue on revenund, Sonochere elenium, Hydronically of low templicon atom; their chemis) to initial silicon-transchemistry	per if necessary and instry, Silinguistry, S	con, Organilylenes, /87 were interested op routes develop to in the all alized	block number) nosilanes, Silacenium
Philip Boudjouk TYPE OF REPORT Final SUPPLEMENTARY NOTATION COSATI CODES FIELD GROUP SUB-GROUP Ultraso Organos ABSTRACT (Continue on reverse if necessary and identify by block The objectives of this program for the 1) to continue to extend our research on ul 2) to continue to develop the new chemistry silvlenes containing large groups on the si silvlenium ions in solution and investigate chemistry of spiropentasilane derivatives; syntheses of novel silicon-chalcogenide and compounds; and, 6) to investigate the elect organosilanes. In this report we summarize meet these objectives.	(Continue on rever und, Sonocher elenium, Hydr number) period 11/1 trasonically to f low temp licon atom; their chemi 5) to initial silicon-tra- rochemistry the results	per if necessary and instry, Silinguistry, S	con, Organilylenes, /87 were direction erated op routes develop to a limite all alized on trying to the second or trying to t	block number) nosilanes, Silacenium
Philip Boudjouk TYPE OF REPORT Final COSATI CODES FIELD GROUP SUB-GROUP The objectives of this program for the l) to continue to extend our research on ul 2) to continue to develop the new chemistry silvlenes containing large groups on the si silvlenium ions in solution and investigate chemistry of spiropentasilane derivatives; syntheses of novel silicon-chalcogenide and compounds; and, 6) to investigate the elect organosilanes. In this report we summarize meet these objectives.	Continue on reversable of low templicon atom; their chemically of low templicon for initial silicon—trachemistry the results	per if necessary and istry, Sili cosilation, S /84 - 10/31/ accelerated erature generature generature; 4) to develor stry; 4) to te programs insition metal of functional obtained in the second of t	con, Organilylenes, /87 were direction erated op routes develop to in the all alized in trying to control organized to the control of the co	block number) nosilanes, Silacenium s; to
Philip Boudjouk The OF REPORT Final FROM 11/1/83 TO 10/31/8 SUPPLEMENTARY NOTATION COSATI CODES 18. SUBJECT TERMS FIELD GROUP SUB-GROUP Ultraso Organos ABSTRACT (Continue on reverse if necessary and identify by block The objectives of this program for the 1) to continue to extend our research on ul 2) to continue to develop the new chemistry silvlenes containing large groups on the si silvlenium ions in solution and investigate chemistry of spiropentasilane derivatives; syntheses of novel silicon-chalcogenide and compounds; and, 6) to investigate the elect organosilanes. In this report we summarize meet these objectives.	Continue on reversable of low templicon atom; their chemically of low templicon for initial silicon—trachemistry the results	per if necessary and istry, Sili cosilation, S /84 - 10/31/ accelerated erature generature generature generature generature generature in the programs in the programs of function and obtained in the control of functions of fu	con, Organilylenes, /87 were reaction erated op routes develop to in the all alized at trying to control trying tryin	block number) nosilanes, Silacenium s; to

"New Approaches to the Synthesis of Novel Organosilanes"

Final Technical Report Submitted to the

Air Force Office of Scientific Research

for Grant No. 84-0008

For the Period November 1, 1983 through October 31, 1987

BY

DR. PHILIP BOUDJOUK PROFESSOR

DEPARTMENT OF CHEMISTRY NORTH DAKOTA STATE UNIVERSITY FARGO, NORTH DAKOTA 58105-5516

TABLE OF CONTENTS

Title	page		• • • • • • •			• • • • • • • • •	1
Table	of Contents		• • • • • • •		• • • • • • • •		ii
Abstra	ct		• • • • • • •				iii
Summar	y of Accomp	lishments	· · · · · · · ·			• • • • • • • •	1
Pertin	ent Researc	h Activities	of the	PI	• • • • • • •		10
Cumula	tive List o	f Publication	ns				13
Person	ne1						18



Agge	ssion For	
DTIC	GRA&1 TAB nounced ification	
i By		
Ava	ribution/ ilability _Avail and	
P	Special	

ABSTRACT OF OBJECTIVES AND ACCOMPLISHMENTS

The objectives of this program for the period 11/1/84 - 10/31/87 were 1) to continue to extend our research on ultrasonically accelerated reactions; 2) to continue to develop the new chemistry of low temperature generated silylenes containing large groups on the silicon atom; 3) to develop routes to silylenium ions in solution and investigate their chemistry; 4) to develop the chemistry of spiropentasilane derivatives; 5) to initiate programs in the syntheses of novel silicon-chalcogenide and silicon-transition metal compounds; and, 6) to investigate the electrochemistry of functionalized organosilanes. In this report we summarize the results obtained in trying to meet these objectives.

Discussion Resistant Resistant Industries

We use ultrasound to prepare the nickel powder from nickel chloride and lithium:

Our examples are the first synthetically successful nickel-catalyzed hydrosilations. The literature on this metal as a catalyst in either heterogeneous or homogeneous form reports very low yields of silicon products. We are investigating the scope of this reaction in detail because of the important economic implications of replacing platinum, which adds more than \$0.12 to cost of each pound of hydrosilated compound, with the much cheaper nickel.

Reductive Silvlation of Carbonyls

In addition to the hydrosilation reaction described above, we discovered that ultrasound will promote the reductive silylation of carbonyl compounds in the presence of zinc and trimethylchlorosilane and will, in some cases, deoxygenate ketones and aldehydes.

$$R_2C=0 + Me_3SiC1 + Zn \xrightarrow{)))} > R_2C=CR_2$$

for benzaldehyde and 1-indanone

The latter reaction holds promise as a new olefin synthesis and is especially attractive because the conditions are mild: room temperature, atmospheric pressure and short reaction times (2-5 hrs). Moreover, the reagents are inexpensive and readily available.

Hydrogenation of Olefins

We have extended our studies of ultrasonically acccelerated reactions to three phase systems. Using a high intensity ultrasonic probe fitted with a stainless steel vessel we have been able to hydrogenate olefins at 0°C in less than 2 hours. The hydrogen is under only 2 atmospheres of pressure!

$$H_2 + C=C$$
 $\xrightarrow{)))}$ $H-C-C-H$ >90% yields

one catalyst is the traditional platinum on carbon with no special preparation and is used in only very small quantities (approximately 0.001 mole percent compared to the quantity of olefin).

Chemistry of Low Temperature Generated Silylenes

Several years ago we reported the room temperature synthesis of d-t-butyl-silylene, a highly hindered silylene with the distinct possibility of forming a triplet ground state. The possibility for this unique ground state arises from the large size of the t-butyl group and the potential that the huge steric requirement might force the divalent intermediate into a linear geometry and therefore a triplet state-all silylenes reported to date have been ground state singlets.

Last year we reported in our annual report the completion of the definitive Skell experiment that demonstrates that this new silvlene is indeed a singlet and that there is little or no triplet present at ambient conditions. Using the sequence of reactions below we have very convincing evidence:

1)
$$t-Bu_2SiCl_2 + Li \xrightarrow{)))$$
 > $t-Bu_2Si: + 2LiCl$

2)
$$t-Bu_2Si: + H C-C H_3 \longrightarrow t-Bu_2Si - C-CH_3 H$$

3)
$$t-Bu_2Si \xrightarrow{C-CH_3} \underbrace{MeO-D}_{C-CH_3} \xrightarrow{MeO-D} t-Bu_2Si \xrightarrow{C-CH_3}_{H}$$

The deuterium-containing carbon in the final product is chiral and its stereochemistry defines the geometry of the silacyclopropane from which it is derived. In summary, the reaction sequence demonstrated that the silacyclopropane was formed by concerted addition to the cis-butene and that there was no stepwise addition to the double bond. Stepwise addition would give some trans dimethyl stereochemistry to the three membered ring and methanolysis would yield a different stereoisomer from the one obtained. Concerted addition is a singlet reaction, stepwise addition is more typical of triplets.

We now wish to report that we have isolated the three membered ring from the reaction of di-t-butylsilylene with cis-butene to obtain unequivocal proof that this silylene is a ground state singlet. There is only one product from this reaction, cis-2,3-dimethyl-1,1-di-t-butylsilacyclopropane, a very stable compound that exhibits unique reactivity in that, despite the absence of chromophores like phenyl or vinyl groups, has an absorption above 254 nm and can be photolyzed. The consequence: we have shown that this silacyclopropane can now liberate di-t-butylsilylene by room temperature photolysis. Alternatively, we can pyrolyze the ring system to give the same species. The net result of all this is that we now have three convenient routes to a reactive intermediate that had been sought after for at least three decades.

These intermediates are very useful for building larger more complex silanes and polysilanes. We have begun investigating the synthetic potential of our method of generating silylenes and we have found that they are indeed very satisfactory as building blocks. Using silylenes generated in this fashion we have been able to prepare novel disilanes and trisilanes as well as some very unusually bulky compounds containing only one silicon atom.


The most significant result of this "building block" is the synthesis of the first stable cyclotetrasilane with functional groups on the silicon atoms:

We have prepared this novel compound in yields as high as thirty percent and we have just completed its crystal structure. The unusual aspect of the structure of this compound is that the hydrogen atoms are trans and <u>perpendicular</u> to the nearly planar ring. Most other rings of this size are puckered.

Silylenium Ions In Solution

We have undertaken an effort to prepare stable silylenium ions in solution

to pursue the possibility that they may serve as useful molten salts in electrochemical applications or as novel initiators for the polymerization of halocarbons. The system currently under investigation is the silole below:



We have completed a series of experiments which indicate that this system may readily produce silylenium ions. For example, hydride abstraction from the silole gives a solution which has a conductance which parallels very closely the conductance of the known organic ionic salt, triphenylcarbenium tetraphenylborate. Stabilization of this silylenium ion is probably from two sources: vinyl stabilization from the two olefinic bonds in the ring system and steric hindrance from the phenyl groups on the ring and on the silicon. We will be examining this system very carefully in the coming year because there are only two examples in the literature of this kind of species. This species has a particularly novel aspect to it: If in fact it is a charge separated silylenium ion then it is the first example of an antiaromatic species containing silicon.

Silicon-Selenium Heterocycles

The silicon-selenium bond is a reactive polarizable bond that has not been examined closely for its potential in synthesis or in the production of novel materials, in particular, non-linear optic materials. We have prepared the new systems below and completed a preliminary examination of their chemistry. In collaboration with Professor Lyle Parker of this Department we have measured the nonlinear optical activity of this ring system and found that the ring with

R - Ph has frequency doubling (from 1060 nm to 532 nm) with 60% the efficiency of urea. This is a very promising result and provides the basis for further exploration of main group elements as skeletal fragments for nonlinear optic materials.

R - Me, Et, Ph

We can at this stage also report a novel reaction: these ring systems readily fracture to give good yields of new reactive intermediates containing the silicon selenium double bond, R₂Si-Se. We are examining the chemistry of this species and will be able for the first time make a comparison with related members of the same family, i.e., compare the chemistry and spectroscopic properties of R₂Si-C, R₂Si-S and R₂Si-Se. Reactive intermediates in which silicon is doubly bonded to another atom have been intensely studied for more than 15 years but only now can a systematic study within a family be executed. We have easy access to all three kinds of species and can compare reaction chemistries. We are collaborating with Hans Bock (Frankfurt) on photoelectron spectroscopic measurements of these reactive intermediates and Dr. Greg Gillispie of our own department to obtain infrared and ultraviolet spectra of these species.

At this point we have demonstrated the existence of this intermediate by trapping reactions with cyclic siloxanes leading to the synthesis of novel ring systems containing silicon, oxygen and selenium.

Electrochemistry

The electrochemistry of functionalized silanes is poorly developed and, as are sult, not well understood. The problems in this potentially important area of chemistry stem from the high sensitivity of halosilanes to water (a persistent contaminant in electrolytes) and the high reduction potentials of these compounds. With considerable effort we were able to obtain the first accurate electrochemical data on a variety of halosilanes (Tables 1 and 2).

Table 1

Reduction Potentials of Some Mono and Dihalosilanes

			
Me ₂ HSiCl	-3.2v	Me ₂ SiCl ₂	-3.2
Me ₃ SiCl	-3.2	t-Bu ₂ SiCl ₂	-3.2
Et ₃ SiCl	-3.2	Ph ₂ SiCl ₂	-3.2
t-BuMe ₂ SiCl	-3.2	$\mathtt{Mes_2SiCl_2}$	-2.8, -3
Ph ₂ HSiCl	-3.2	(t-Bu ₂ Si) ₂ Br	-2.9, -3
Ph ₃ SiCl	-2.85	Dt	

Table 2

Reduction Potentials of Functionalized Silanes

Me ₃ Si-OTs	-2.9 v
Me ₃ Si-OCPh ₃	-3.2
Me ₃ Si-OPh	-3.2
Et3Si-OAc	-3.2
Ph ₂ HSi-OPh	-3.2

Moreover, we were able to translate these data to a workable synthetic scheme and produce disilanes in high yields. The approach is quite limited in the laboratory because of the care required for each experiment. There is more cause for optimism on very large scale, however, because the system is amenable to flow synthesis, a scheme that eliminates many of the problems incurred in batch synthesis. Table 3 summarizes some of our results:

Table 3

Electrosynthesis of Some Disilanes

Me ₂ SiHCl	HMe ₂ Si-SiMe ₂ H	88	82
Me ₃ SiCl	Me ₃ Si-SiMe ₃	76	60
Et ₃ SiCl	Et ₃ Si-SiEt ₃	77	67
CH2=CHMe2SiCl	CH2-CHMeSi)2	83	73
Ph ₂ HSiCl	Ph ₂ HSi-SiHPh	90	70

References

- 1. B.-H. Han and P. Boudjouk, Organometallics 1983, 2, 769-771.
- P. Boudjouk, XVIIIth National Organosilicon Symposium, Schenectady, NY April 1984.
- 3. J. Lambert, and W. Schulz, Jr., . J. Am. Chem. Soc. 1983, 105, 1671-1672;
 - J. Lambert, JoAnn McConnell and W. Schulz, Jr., J. Am. Chem. Soc., 1986, 108, 2482-2484.

Pertinent Recent Research Activities of the Principal Investigator

The research supported by AFOSR through grants 80-0239 and 84-0008 has led to over twenty five papers (including two reviews) and three book chapters. The thrust of our work has been in organosilicon chemistry but, as a result of our efforts to improve the reaction rates of organosilanes, we developed a program to investigate the effects of ultrasonic waves in synthesis. This relatively new area is now known as "sonochemistry" and it is growing rapidly. The AFOSR is well-known as one of the first to support research in this field.

That the interest in sonochemistry is escalating seems clear.

- 1) Symposia: two symposia have already been held that focussed on the chemical effects of ultrasonic waves, the first at the University of Warwick, England, April 1986 as part of the Annual Chemical Congress of the Royal Society of Chemistry. A second was held as part of the ACS Meeting in Anaheim, California, September 1986. The PI was an invited speaker for both symposia.
- 2) Research papers: while it is difficult to arrive at an exact figure for the the number of papers dealing with ultrasonic waves in synthesis, the recent activity in sonochemistry have justified several reviews. Suslick has a major review on mostly homogeneous sonochemistry and the PI has written three on heterogeneous sonochemistry, the most recent has 30 tables and over 250 references dealing mostly with heterogeneous applications. Additionally, numerous reviews have been published in Great Britain, France, Japan and Russia.
- 3) Industrial interest: since our first paper in 1981, industrial interest has been high. Over two-hundred companies have called and written to inquire about somochemistry and the PI has presented seminars on this topic at about forty

companies in the last four years. Several companies report excellent success on reactions of interest to them but have had difficulty scaling up to more than 25 gallons. Our own work has led to two patents.

Research into the effects of ultrasonic waves required that we investigate many of the traditional organic reactions in addition to those involving silicon. The breadth of our coverage is indicated by the titles of the papers in the Publication List but, in summary, we have demonstrated that nearly every kind of heterogenous reaction can be significantly enhanced when sonicated. By "enhanced" we mean increases in rate or yield and nearly all of the time, both. Reactions involving metals (catalytic and stoichiometric) respond most favorably to ultrasound but solids such as potassium hydride, lithium aluminum hydride, potassium permanganate and aluminum oxide are also activated. Supported catalysts also have greater activity when sonicated.

The nature of cavitation is such that greater reaction rates are possible at lower temperatures. This is mainly the result of reduced vapor pressure in the cavity as it collapses giving rise to a more energetic shook wave. The consequence of this is that we have observed the some reactions progress at low temperatures more efficiently than at higher temperatures. For example, hydrogenation of olefins using hydrogen gas and a commercially available Pt/C catalyst is faster and gives better yields at 0° with ultrasound than at room temperature with vigorous mixing[41]. Several Friedel-Crafts and Diels-Alder reactions respond in the same fashion[52].

The successes in synthesis have begun to attract chemists and physicists to the area that were not formerly involved in ultrasound research. There have been numerous studies on cavitation and related phenomena on very ideal systems, but now that "useful" reactions have benefitted from these phenomena there is a

growing interest to reexamine ultrasound under "real" reaction conditions, i.e., nonaqueous solvents, with organic halides present, with metals under normal reaction conditions rather than highly purified and with conventional ultrasonic equipment. Our plan is get involved in these investigations in a collaborative role until we acquire the expertise to conduct our own studies.

PUBLICATIONS

- 1. "Tin Heterocycles. Some Phenoxastannin and Phenothiastannin Derivatives," E.J. Kupchick, J.A. Ursino, and P. Boudjouk, J. Organometal Chem., 10, (1967) 269.
- 2. "New Anionic Rearrangements. IX. 1,2-Anionic Rearrangement of Organosilylhydroxylamines," R. West, P. Boudjouk, A. Matuszko, J. Amer. Chem. Soc., 91, (1969) 5184.
- 3. "A Novel Thermal Rearrangement of Tris(organosily1)hydroxylamines," P. Boudjouk and R. West, J. Am. Chem. Soc., 93, (1971) 5901.
- 4. "Bis(organosily1)nitroxides," R. West and P. Boudjouk, J. Amer. Chem. Soc., 93, (1971) 5902.
- 5. "New Anionic Rearrangements, XVI. 1, 2-Anionic Rearrangement from Sulfur to Carbon in Benzylthiotrimethylsilane," A. Wright, D. Ling, P. Boudjouk, and R. West, J. Amer. Chem. Soc., 94, (1972) 4784.
- 6. "Photochemical Dehydrosilylation of Pentaphenylmethyldisilane. Generation and Trapping of an Unstable Intermediate Containing a Silicon-Carbon Double Bond or Its Equivalent." P. Boudjouk, J.R.Roberts, C. M. Golino, and L.H. Sommer, J. Amer. Chem. Soc., 92, (1972) 7926.
- 7. "Photochemical Generation of an Intermediate Containing a Silicon-Carbon Double Bond or Its Equivalent from 1,1-Diphenylsilacyclobutane," P. Boudjouk and L.H. Sommer, Chem. Commun., (1973) 54.
- "Organosilyl and Organogermyl Nitroxides. A New Radical Rearrangement,"
 R. West and P. Boudjouk, J. Amer. Chem. Soc., 95, (1973) 3983.
- 9. "New Anionic Rearrangements. XV. 1,2-Anionic Rearrangement of Organosilylhydroxylamines," R. West and P. Boudjouk, J. Amer. Chem. Soc., 95, (1973) 3987.
- 10. "Organosilyhydroxylamines," P. Boudjouk and R. West, Intra-Sci. Chem. Rpt., 7, (1973) 65.
- 11. "The Thermal Disproportionation of 3-Chloropropyltrichlorogermane," P. Boudjouk, Inor. Nucl. Chem. Letters, 11, (1975) 679.
- 12. "Vacuum-Ultraviolet Photolysis of C₂(CH₃)₆, Si₂(CH₃)₆ and (CH₃)₃CSi(CH₃)₃. Evidence for an Unsaturated Si-C Linkage," P. Boudjouk and R.D. Koob, J. Amer. Chem. Soc., <u>97</u>, (1975) 6595.
- 13. "Irreversible Thermal Rearrangment of Tris(organosilyl)hydroxylamines," R. West, P. Nowakowski and P. Boudjouk, J. Amer. Chem. Soc., 98, (1976) 5620.

- 14. "The Molecular Structure of 1-Methyl-1-Silabicyclo[2.2.1]heptane by Gas Phase Electron Diffraction. Structural Support for the Trigonal Bipyramidal Transition State in S_N2-Si Reactions," R.L. Hilderbrandt, G.D. Homer and P. Boudjouk, J. Amer. Chem. Soc., 98, (1976) 7476.
- 15. "A Convenient Synthesis of 1-Bromo-8-iodonaphthalene and 1,8-Dibromo-naphthalene from 8-Bromo-1-naphthoic Acid", J.S. Kiely, L.L. Nelson and P. Boudjouk, J. Org. Chem., 42, (1977) 1480.
- 16. "A Synthesis of Terminal Arylacetylenes An in situ Generated Copper (I) Acetylide", J.S. Kiely, P. Boudjouk and L.L. Nelson, J. Org. Chem., 42 (1977) 2626.
- 17. "The Synthesis of 1-Methyl-1-Silaadamantane", C.A. Kapfer and P. Boudjouk, J. Organometal. Chem., 144, (1978) C6.
- 18. "The Photolysis of 1,1-Dimethylsilacyclobutane", R.D. Koob, P. Boudjouk, and S. Tokach, J. Phys. Chem., 82, (1978) 1203.
- 19. "Vinyl-Cylcohexatriene Iron Tricarbonyl Complexes of the Diene Type via Olefin Isomerization of Allylaromatics", P. Boudjouk and S. Lin. J. Organometal. Chem., 155, (1978) C13.
- 20. "Improved Routes to Phenalene and Phenalanone. Alane, Borane, and Silane Reductions of Phenalenone", P. Boudjouk and P.D. Johnson, J. Org. Chem., 43, (1978) 3979.
- 21. "The Structure of 1-Methyl-1-Silaadamantane as Determined by Gas Phase Electron Diffraction", Q. Shen, C.A. Kapfer, P. Boudjouk, and R.L. Hilderbrandt, J. Organometal. Chem., 169, (1979) 147.
- 22. "The Organometallic Chemistry of Phenalene. Neutral and Cationic n² Complexes Phenalene", J. Woell and P. Boudjouk, J. Organometal. Chem., <u>172</u>, (1979) C43.
- 23. "The Structure of 1-Methyl-1-Germandamantane as Determined by Gas Phase Electron Diffraction", Q. Shen, C.A. Kapfer, P. Boudjouk, and R.L. Hilderbrandt, J. Molec. Struct., <u>54</u>, (1979) 295.
- 24. "1,8-Difunctional Naphthalenes as Building Blocks. A Convenient Low Temperature Synthesis of Silaacenaphthenes", J.S. Kiely and P. Boudjouk, J. Organometal. Chem., 182, (1979) 173.
- 25. "On the Purported Photochemical Oxidation of Alcohols by Ag(I). A Re-examination", R.S. Macomber, S. Carr, P. Boudjouk, and C.A. Kapfer, J. Org. Chem., 45, (1980) 356.
- 26. "The Organometallic Chemistry of Phenalene. Neutral n^1 , n^3 , n^6 Complexes of Phenalene", S. Lin and P. Boudjouk, J. Organometal. Chem., 187. (1980) C11.

- 27. "A Convenient and Unambiguous Synthesis of 1-Bromoindene", J.B. Woell and P. Boudjouk, J. Org. Chem., 45, (1980) 5213.
- 28. "The Reaction of Isobutylene Dicarbonylcyclopentadienyliron Tetrafluoro-borate with Diphenylcyclopropenone: Complexation Without Ring Gpening", J.B. Woell and P. Boudjouk, Angew. Chem. Internat. Edit. 20, (1981) 387.
- 29. "Organic Sonochemistry. Ultrasound-Promoted Coupling of Organic Halides in the Presence of Lithium Wire", B-H. Han and P. Boudjouk, Tetrahedron Lett., 2, (1981) 2757.
- 30. "Organic Sonochemistry. Ultrasound Promoted Coupling of Chlorosilanes in the Presence of Lithium Wire", P. Boudjouk and B-H. Han, Tetrahedron Lett., 22, (1981) 3813.
- 31. "The Synthesis of 1-Silaphenalenes from 1-8-Difunctional Naphthalenes. Confirmation of the Structure of the Pyrolysis Products of (1-Naphthyl)-vinyldichlorosilane", P. Boudjouk, J.S. Kiely and R. Sooriyakumaran, J. Organometal. Chem., 221 (1981) 33.
- 32. "The Synthesis of the Elusive 1,8-Bis(trimethylsilyl)naphthalene and Its Facile Rearrangement to the 1,7 Isomer", R. Sooriyakumaran and P. Boudjouk, Organometallics, 1, (1982) 218.
- 33. "Organotransition Metal Complexes of Very Basic Ketones. The Synthesis of Dicarbonylcyclopentadienyliron Complexes of Diphenylcyclopropenone, Tropone and Phenalenone. The Crystal and Molecular Structure of Dicarbonylcyclopentadienyliron Tropone Tetrafluoroborate", P. Boudjouk, J.B. Woell, L.J. Radonovich and M.W. Eyring, Organometallics, 1 (1982) 582.
- 34. "Organic Sonochemistry. Ultrasound Promoted Reaction of Zinc with alpha, alpha-Dibromo-o-xylene. Evidence for Facile Generation of o-Xylylene", B-H. Han and P. Boudjouk, J. Org. Chem., <u>47</u> (1982) 751.
- 35. "Organic Sonochemistry. Ultrasound Acceleration of the Reduction of Simple and Deactivated Aryl Halides Using Lithium Aluminum Hydride", B-H. Han and P. Boudjouk, Tetrahedron Letters, 23, (1982) 1643.
- 36. "Organic Sonochemistry. Sonic Acceleration of the Reformatsky Reaction", B-H. Han and P. Boudjouk, J. Org. Chem., 47, (1982) 5030.
- 37. "Sonochemical and Electrochemical Synthesis of Tetramesityldisilene", P. Boudjouk, B-H. Han and K.R. Anderson, J.Amer.Chem. Soc., <u>104</u>, (1982) 4992.
- 38. "Palladium-Catalyzed and Sonically Accelerated Hydrogenations of Olefins Using Formic Acid as a Hydrogen Transfer Agent", P. Boudjouk and B-H. Han, J. Catalysis, 79(1983) 489.
- 39. "The Synthesis and Reactivity of 1-Silaadamantyl Systems," P. Boudjouk, C.A. Kapfer, and R.F. Cunico, Organometallics, 2 (1983) 336.

- 40. "Organic Sonochemistry. Ultrasonic Acceleration of the Hydrosilation Reaction", B-H. Han and P. Boudjouk, Organometallics, 2, (1983) 769.
- 41. "Are the Silacyclopentadienyl Anion and the Silacyclopropenyl Cation Aromatic?" H.S. Gordon, P. Boudjouk and F. Anwari, J. Amer. Chem. Soc., 105, (1983) 4972.
- 42. "Organische Chemie mit Ultraschall", P. Boudjouk, NACHRICHTEN aus Chemie, Technik and Laboratorium, 31, (1983) 798.
- 43. "Theoretical Studies of Polyvinyl-Substituted Carbenium and Silylenium Ions" T. Truong, M.S. Gordon and P. Boudjouk, Organometallics, 3, (1984) 484.
- 44. "1-Silaphenalenes, Potential Precursors to Aromatic Silylenium Ions and Aromatic Silyl Anions. Spectroscopic and Chemical Studies", R. Sooriya-kumaran and P. Boudjouk, J. Organometal. Chem. 271, (1984) 289.
- 45. "The Synthesis of the First Spiropentasilane, Octamethylspiropentasilane" P. Boudjouk and R. Sooriyakumaran, J.Chem. Soc., Chem. Commun. (1984)777.

Banda Sales

- 46. "The Reaction of Magnesium with cis-1,3,5-Tris(bromomethyl)cyclohexane. Evidence for a Soluble Tri-Grignard", P. Boudjouκ, R. Sooriyakumaran and C.A. Kapfer, J. Organometal. Chem., 281, (1985) C21.
- 47. "The Molecular and Electronic Structures of Metallospiropentanes" M.S. Gordon and P. Boudjouk, J. Amer. Chem. Soc., 107, (1985) 1439.
- 48. "Structure of 1,8-Bis(trimethylsilyl)naphthalene", R. Sooriyakumaran, P. Boudjouk and R.G. Garvey, Acta Crystallagraphia C41, (1985) 1348
- 49. "The Synthesis of 1-Methyl-1-Germaadamantane", P. Boudjouk and C.A. Kapfer, J. Organometal. Chem., <u>296</u>, (1985) 339.
- 50. "An Improved Synthesis of Hexamethylphenalene", P. Boudjouk, W.H. Ohrbom and J.B. Woell, Synthetic Commun., 16, 401(1986).
- 51. "Synthesis with Ultrasonic Waves", P. Boudjouk, J. Chem. Ed., 63, (1986) 427.
- 52. "Organic Sonochemistry. Ultrasonic Acceleration of the Reaction of Dicarbonyls with Trimethylchlorosilane in the Presence of Zinc", P. Boudjouk and J.H. So, Synth. Commun., 16, (1986) 775.
- 53. "Ultrasonic Generation of Metal Powders from Lithium and Metal Halides" Philip Boudjouk, Dennis Thompson, Walter H. Ohrbom and Byung-Hee Han, Organometallics, 5, (1986) 1257
- 54. "Organic Sonochemistry. New Sonically Accelerated Reactions Involving Lithium" P. Boudjouk, R. Sooriyakumaran and B.-H. Han, J. Org. Chem., 51, (1986) 2818.

- 55. "Generation and Trapping of Diethylsilaselanone. A Reactive Intermediate with a Silicon-Selenium Double Bond." D. P. Thompson and P. Boudjouk, J. Chem. Soc. Chem. Commun., 1466, 1987.
- 56. "A Convenient, Naphthalene-Catalyzed Synthesis of Alkali Metal Selenides and Diselenides in Tetrahydrofuran and the Reactivity Differences Exhibited by These Salts Toward Organic Bromides. Effect of Ultrasound.", D. P. Thompson and P. Boudjouk, J. Org. Chem., in press.
- 57. "Ultrasonically Accelerated Nickel Catalyzed Hydrosilation Reactions", P. Boudjouk, B.-H. Han and J. Jacobsen, submitted.
- 58. "Ultrasonic Acceleration of the Reaction of Lithium with Bulky Diorganodichlorodisilanes. Generation and Trapping of Di-t-Butylsilylene, a Ground State Triplet." P. Boudjouk, K.R. Anderson, R. Sooriyakumaran, U. Samaraweera and J. Chrusciel, submitted
- 59. "A Convenient Synthesis of Binary and Ternary Selenides", D. P. Thompson and P. Boudjouk, submitted.
- 60. "An Efficient Route to Unsymmetrical Granoselenides", D. P. Thompson and P. Boudjouk, submitted.
- 61. "Sonochemistry", K. S. Suslick and P. Boudjouk, Acc. Chem. Res., submitted.

Book Chapters

- 1. "Electrochemical and Sonochemical Routes to Organosilane Precursors", in <u>Science of Ceramic Chemical Processing</u>, L.L. Hench and D.R. Ulrich, Eds., John Wiley & Sons, Inc., New York, 1986
- "The Acceleration of Synthetically Useful Heterogeneous Reactions Using Ultrasonic Waves.", P. Boudjouk, in "High Energy Processes in Organometallic Chemistry", K. S. Suslick, Ed., American Chemical Society Symposium Series, American Chemical Society, Wash. DC, 1987.
- 3. "Heterogeneous Sonochemistry", in <u>Ultrasound: Chemical</u>, <u>Physical and Biological Effects</u>, K.S. Suslick, Ed., Verlag Chemie International, in press.

Personnel

Senior Personnel

Philip Boudjouk, Principal Investigator

Byung-Hee Han, Research Associate and Visiting Professor Moon Kyeu Park, Research Associate and Visiting Professor

Jerzy Chrusciel, Research Associate Yogendra Gupta, Research Associate Steven Keller, Research Associate

Junior Personnel (Graduate Students)

Kevin Anderson
Steven Bahr
Amirthini Balasingham
Eric Black
Elizabeth Bonderson
John Jacobsen
Andrew King
Rajkumar Kumarathasan
Walter Ohrbom
Upasiri Samaraweera
Jeung Ho So
Ratnasabapathy Sooriyakumaran
Dennis Thompson

SES TESTICAL CASSING TO SESSION